

# IMPACTS OF SEDIMENT ON KŌARO



Sediment can affect māhinga kai by influencing habitat, behaviour, feeding, growth and survival.

## Background on kōaro (*Galaxias brevipinnis*)

Kōaro are one of six species in Aotearoa New Zealand's whitebait catch. They are diadromous – they spend over 4½ months in the ocean as larvae before entering rivers<sup>1</sup>. Kōaro make up about 5% of the whitebait caught in rivers around the whole of New Zealand, but they are rare in east coast regions, particularly Canterbury<sup>2</sup>. Kōaro whitebait mostly migrate into cold water rivers coming from glaciers and mountain regions<sup>3</sup>. Seagoing populations can migrate large distances inland (250 km upstream) and to over 1,000 m of altitude<sup>4</sup>. In addition to seagoing populations, kōaro also form lake-limited populations in which the larvae do not go to sea but develop in lakes<sup>5</sup>. Kōaro reach sexual maturity after two years<sup>6</sup> and almost certainly spawn several times. They probably live for six to eight years, perhaps longer<sup>7</sup>. Kōaro are the only New Zealand freshwater fish found on the sub-Antarctic islands of New Zealand and are found also in Tasmania and mainland Australia<sup>7</sup>.

## Kōaro (*Galaxias brevipinnis*)



## Kōaro sensitivity to elevated sediment



■ Low   ■ Medium   ■ High

Prepared by Mike Hickford, Michele Melchior and Melanie Mayall-Nahi from NIWA for Our Land and Water National Science Challenge, March 2023. Image of kōaro by Dr R M McDowall.

For references and further information see [niwa.co.nz/sediment-impacts](https://niwa.co.nz/sediment-impacts)

# IMPACTS OF SEDIMENT ON KŌARO

## Effects of suspended sediment on kōaro

|                  |  |
|------------------|--|
| <b>Habitat</b>   | Adult kōaro are found in rapidly flowing, tumbling rocky streams in native forest <sup>7</sup> . Studies in West Coast rivers show kōaro are abundant only in clear-flowing rocky streams, with few occurring in brown, tannin-stained streams. Kōaro are rarely found outside forests, and then, only in streams that have just left forest-cover or in the tributaries of high-country lakes. However, kōaro appear relatively insensitive to increases in suspended sediments in rivers <sup>4</sup> and their occurrence is not related to the duration of turbid conditions in rivers <sup>4</sup> .                      |
| <b>Behaviour</b> | In the laboratory, kōaro whitebait quickly move from highly turbid water to less turbid water <sup>8</sup> . However, kōaro whitebait are relatively more abundant in snow-fed rivers, which tend to become turbid during the spring snow thaw, than in rivers draining coastal hills <sup>4,9</sup> . This is not to say that kōaro whitebait prefer turbid waters, just that the proportion of kōaro whitebait in catches is often higher immediately after floods, when the water is still turbid <sup>9</sup> .  |
| <b>Feeding</b>   | Kōaro whitebait are better able to feed in turbid waters than the other whitebait species. Even very high turbidity levels have no significant effect on the mean feeding rate of juvenile kōaro <sup>10</sup> . This is despite vision being important for feeding by kōaro whitebait - the absence of light reduces their feeding by 90% <sup>10</sup> . Although high turbidity does not affect the feeding rate of kōaro juveniles, even short-term exposure (two hours) to high turbidity levels stresses them such that when they return to clear water their feeding rates are suppressed significantly <sup>10</sup> . |
| <b>Growth</b>    | Sustained periods (21 days) of moderate levels of turbidity reduce the growth (length, but not weight) of juvenile kōaro <sup>11</sup> . This is probably an indirect effect caused by reduced feeding efficiency.   |
| <b>Survival</b>  | Long-term (21 days) exposure to very high turbidity has no effect on the survival of juvenile kōaro <sup>11</sup> .  |

## Effects of deposited sediment on kōaro

|                  |   |
|------------------|---|
| <b>Habitat</b>   | <p>Kōaro usually live in the swifter rapids of streams, under logs, or more usually, under and amongst boulders<sup>7</sup>. During the day and night, kōaro use microhabitats that are faster-flowing, deeper, and more turbulent, with larger substrate particles and more interstitial (between substrate particles) refuge spaces than elsewhere in the reach<sup>12,13</sup>. Interstitial spaces are important habitat for kōaro, so deposited sediments will reduce the amount of interstitial space available in streams and cause population decline<sup>14</sup>.</p> <p>It is likely that kōaro spawn high on stream banks during flood events and that their eggs are deposited well above baseflow water levels<sup>15,16</sup>. The few records of kōaro spawning sites in New Zealand<sup>15</sup> and Australia<sup>17</sup> show eggs are laid in the interstices of cobble and pebble substrate adjacent to the stream channel even though other sediment types are more commonly available. Any sediment that is deposited on stream margins during floods will reduce the availability of the preferred spawning substrate.</p> |
| <b>Behaviour</b> | Direct effects unknown.   |
| <b>Feeding</b>   | Benthic invertebrates are numerically the greatest food source for kōaro, but larger, terrestrial prey are more important by weight <sup>18</sup> . High sediment loads can affect benthic invertebrate communities, generally causing increased drift, along with reduced densities and biomass <sup>19-21</sup> . It is possible that deposited sediments may reduce some key items in the diet of kōaro.   |
| <b>Growth</b>    | Direct effects unknown.   |
| <b>Survival</b>  | Direct effects unknown.   |

# IMPACTS OF SEDIMENT ON KŌARO

## Further information:

1. McClintock, G.J. (2018). Early life history dynamics of the New Zealand whitebait species. M.Sc. thesis. University of Canterbury: Christchurch. 106p.
2. Yungnickel, M.R., M.J.H. Hickford, and D.R. Schiel (2020). Spatio-temporal variation in species composition of New Zealand's whitebait fishery. *New Zealand Journal of Marine and Freshwater Research* 54(4): 679-694.
3. McDowall, R.M. (1965). The composition of the New Zealand whitebait catch, 1964. *New Zealand Journal of Science* 8(3): 285-300.
4. Rowe, D.K., D.M. Hicks, and J. Richardson (2000). Reduced abundance of banded kokopu (*Galaxias fasciatus*) and other native fish in turbid rivers of the North Island of New Zealand. *New Zealand Journal of Marine and Freshwater Research* 34(3): 547-558.
5. Rowe, D.K., G. Konui, and K.D. Christie (2002). Population structure, distribution, reproduction, diet, and relative abundance of koaro (*Galaxias brevipinnis*) in a New Zealand lake. *Journal of the Royal Society of New Zealand* 32(2): 275-291.
6. Eldon, G.A. (1969). Observations on growth and behaviour of Galaxiidae in aquariums. *Tuatara* 17(1): 34-46.
7. McDowall, R.M. (1990). *New Zealand freshwater fishes: a natural history and guide*. Auckland: Heinemann Reed. 553p.
8. Boubée, J.A.T., T.L. Dean, D.W. West, and R.F.G. Barrier (1997). Avoidance of suspended sediment by the juvenile migratory stage of six New Zealand native fish species. *New Zealand Journal of Marine and Freshwater Research* 31(1): 61-69.
9. McDowall, R.M. and G.A. Eldon (1980). The ecology of whitebait migrations (*Galaxiidae: Galaxias* spp.). *New Zealand Ministry of Agriculture and Fisheries, Fisheries Research Bulletin* 20: 1-172.
10. Rowe, D.K. and T.L. Dean (1998). Effects of turbidity on the feeding ability of the juvenile migrant stage of six New Zealand freshwater fish species. *New Zealand Journal of Marine and Freshwater Research* 32(1): 21-29.
11. Cavanagh, J.E., K.L. Hogsdon, and J.S. Harding (2014). Effects of suspended sediment on freshwater fish, in Landcare Research Contract Report No. LC1986, 2p.
12. McEwan, A.J. (2009). Fine scale spatial behaviour of indigenous riverine fish in a small New Zealand stream. M.Sc. thesis. Massey University: Palmerston North. 95p.
13. McEwan, A.J. and M.K. Joy (2014). Diel habitat use of two sympatric galaxiid fishes (*Galaxias brevipinnis* and *G. postvectis*) at two spatial scales in a small upland stream in Manawatu, New Zealand. *Environmental Biology of Fishes* 97(8): 897-907.
14. Richardson, J. and I. Jowett (2002). Effects of sediment on fish communities in East Cape streams, North Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 36(2): 431-442.
15. Allibone, R.M. and D. Caskey (2000). Timing and habitat of koaro (*Galaxias brevipinnis*) spawning in streams draining Mt Taranaki, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 34(4): 593-595.
16. Charteris, S.C. (2002). Spawning, egg development and recruitment of diadromous galaxiids in Taranaki, New Zealand. M.Sc. thesis. Massey University: Palmerston North. 89p.
17. O'Connor, W.G. and J.D. Koehn (1998). Spawning of the broad-finned *Galaxias*, *Galaxias brevipinnis* Günther (*Pisces: Galaxiidae*) in coastal streams of southeastern Australia. *Ecology of Freshwater Fish* 7(2): 95-100.
18. Main, M.R. (1988). Factors influencing the distribution of kokopu and koaro (*Pisces: Galaxiidae*). M.Sc. thesis. University of Canterbury: Christchurch. 127p.
19. Newcombe, C.P. and D.D. MacDonald (1991). Effects of suspended sediments on aquatic ecosystems. *North American Journal of Fisheries Management* 11(1): 72-82.
20. Birtwell, I.K. (1999). The effects of sediment on fish and their habitat, in Canadian Stock Assessment Secretariat Report No. 99/139. Department of Fisheries and Oceans Canada: Ottawa 34p.
21. Greenwood, M.J. and D.J. Booker (2016). Influence of hydrological regime and land cover on traits and potential export capacity of adult aquatic insects from river channels. *Oecologia* 180(2): 551-566.

*Prepared by Mike Hickford, Michele Melchior and Melanie Mayall-Nahi from NIWA for Our Land and Water National Science Challenge, March 2023. Image of kōaro by Dr R M McDowall.*